

2009 H1N1 INFLUENZA VACCINE

INACTIVATED (the “flu shot”)

WHAT YOU NEED TO KNOW

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis.

1 What is 2009 H1N1 influenza?

2009 H1N1 influenza (also called Swine Flu) is caused by a new strain of influenza virus. It has spread to many countries.

Like other flu viruses, 2009 H1N1 spreads from person to person through coughing, sneezing, and sometimes through touching objects contaminated with the virus.

Signs of 2009 H1N1 can include:

- Fatigue
- Fever
- Sore Throat
- Muscle Aches
- Chills
- Coughing
- Sneezing

Some people also have diarrhea and vomiting.

Most people feel better within a week. But some people get pneumonia or other serious illnesses. Some people have to be hospitalized and some die.

2 How is 2009 H1N1 different from regular (seasonal) flu?

Seasonal flu viruses change from year to year, but they are closely related to each other.

People who have had flu infections in the past usually have some immunity to seasonal flu viruses (their bodies have built up some ability to fight off the viruses).

The 2009 H1N1 flu is a new flu virus. It is very different from seasonal flu viruses.

Most people have little or no immunity to 2009 H1N1 flu (their bodies are not prepared to fight off the virus).

3 2009 H1N1 influenza vaccine

Vaccines are available to protect against 2009 H1N1 influenza.

- These vaccines are made just like seasonal flu vaccines.
- They are expected to be as safe and effective as seasonal flu vaccines.
- They will not prevent “influenza-like” illnesses caused by other viruses.
- They will not prevent seasonal flu. *You should also get seasonal influenza vaccine, if you want to be protected against seasonal flu.*

Inactivated vaccine (vaccine that has killed virus in it) is injected into the muscle, like the annual flu shot. **This sheet describes the inactivated vaccine.**

A **live, intranasal** vaccine (the nasal spray vaccine) is also available. It is described in a separate sheet.

Some inactivated 2009 H1N1 vaccine contains a preservative called thimerosal to keep it free from germs. Some people have suggested that thimerosal might be related to autism. In 2004 a group of experts at the Institute of Medicine reviewed many studies looking into this theory, and found no association between thimerosal and autism. Additional studies since then reached the same conclusion.

4 Who should get 2009 H1N1 influenza vaccine and when?

WHO

Groups recommended to receive 2009 H1N1 vaccine first are:

- Pregnant women
- People who live with or care for infants younger than 6 months of age
- Health care and emergency medical personnel
- Anyone from 6 months through 24 years of age
- Anyone from 25 through 64 years of age with certain chronic medical conditions or a weakened immune system

As more vaccine becomes available, these groups should also be vaccinated:

- Healthy 25 through 64 year olds
- Adults 65 years and older

The Federal government is providing this vaccine for receipt on a voluntary basis. However, state law or employers may require vaccination for certain persons.

WHEN

Get vaccinated as soon as the vaccine is available.

Children through 9 years of age should get **two doses** of vaccine, about a month apart. Older children and adults need only one dose.

5 Some people should not get the vaccine or should wait

You should not get 2009 H1N1 flu vaccine if you have a **severe (life-threatening) allergy to eggs**, or to **any other substance in the vaccine**. *Tell the person giving you the vaccine if you have any severe allergies.*

Also tell them if you have ever had:

- a life-threatening allergic reaction after a dose of seasonal flu vaccine,
- Guillain Barré Syndrome (a severe paralytic illness also called GBS).

These may not be reasons to avoid the vaccine, but the medical staff can help you decide.

If you are moderately or severely ill, you might be advised to wait until you recover before getting the vaccine. If you have a mild cold or other illness, there is usually no need to wait.

Pregnant or breastfeeding women can get inactivated 2009 H1N1 flu vaccine.

Inactivated 2009 H1N1 vaccine may be given at the same time as other vaccines, including seasonal influenza vaccine.

6 What are the risks from 2009 H1N1 influenza vaccine?

A vaccine, like any medicine, could cause a serious problem, such as a severe allergic reaction. But the risk of any vaccine causing serious harm, or death, is extremely small.

The virus in inactivated 2009 H1N1 vaccine has been killed, so you cannot get influenza from the vaccine.

The risks from inactivated 2009 H1N1 vaccine are similar to those from seasonal inactivated flu vaccine:

Mild problems:

- soreness, redness, tenderness, or swelling where the shot was given
- fainting (mainly adolescents)
- headache, muscle aches
- fever
- nausea

If these problems occur, they usually begin soon after the shot and last 1-2 days.

Severe problems:

- Life-threatening allergic reactions to vaccines are very rare. If they do occur, it is usually within a few minutes to a few hours after the shot.
- In 1976, an earlier type of swine flu vaccine was associated with cases of Guillain-Barré Syndrome (GBS). Since then, flu vaccines have not been clearly linked to GBS.

7 What if there is a severe reaction?

What should I look for?

Any unusual condition, such as a high fever or behavior changes. Signs of a severe allergic reaction can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat or dizziness.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** the doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your provider to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form. Or you can file this report through the VAERS website at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not provide medical advice.

8 Vaccine injury compensation

If you or your child has a reaction to the vaccine, your ability to sue is limited by law.

However, a federal program has been created to help pay for the medical care and other specific expenses of certain persons who have a serious reaction to this vaccine. For more information about this program, call **1-888-275-4772** or visit the program's website at: www.hrsa.gov/countermeasurescomp/default.htm.

9 How can I learn more?

- Ask your provider. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/h1n1flu or www.cdc.gov/flu
- Visit the web at www.flu.gov



DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION



INACTIVATED INFLUENZA VACCINE

WHAT YOU NEED TO KNOW 2009-10

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis.

1 Why get vaccinated?

Influenza (“flu”) is a contagious disease.

It is caused by the influenza virus, which can be spread by coughing, sneezing, or nasal secretions.

Other illnesses can have the same symptoms and are often mistaken for influenza. But only an illness caused by the influenza virus is really influenza.

Anyone can get influenza, but rates of infection are highest among children. For most people, it lasts only a few days. It can cause:

- fever
- sore throat
- chills
- fatigue
- cough
- headache
- muscle aches

Some people, such as infants, elderly, and those with certain health conditions, can get much sicker. Flu can cause high fever and pneumonia, and make existing medical conditions worse. It can cause diarrhea and seizures in children. On average, 226,000 people are hospitalized every year because of influenza and 36,000 die – mostly elderly. **Influenza vaccine can prevent influenza.**

2 Inactivated influenza vaccine

There are two types of seasonal influenza vaccine:

1. **Inactivated** (killed) vaccine, or the “flu shot” is given by injection into the muscle. 2. **Live, attenuated** (weakened) influenza vaccine is sprayed into the nostrils. *This vaccine is described in a separate Vaccine Information Statement.*

These “seasonal” influenza vaccines are formulated to prevent annual flu. They do not protect against pandemic H1N1 influenza.

Influenza viruses are always changing. Because of this, influenza vaccines are updated every year, and an annual vaccination is recommended.

Each year scientists try to match the viruses in the vaccine to those most likely to cause flu that year. When there is a close match the vaccine protects most people from serious influenza-related illness. But even when there is not a close match, the vaccine provides some protection. Influenza vaccine will *not* prevent “influenza-like” illnesses caused by other viruses.

It takes up to 2 weeks for protection to develop after the shot. Protection lasts up to a year.

Some inactivated influenza vaccine contains a preservative called thimerosal. Some people have suggested that thimerosal may be related to developmental problems in children. In 2004 the Institute of Medicine reviewed many studies looking into this theory and concluded that there is no evidence of such a relationship. Thimerosal-free influenza vaccine is available.

3 Who should get inactivated influenza vaccine?

Anyone who wants to **reduce the likelihood of becoming ill with influenza or spreading influenza to others.**

All children 6 months and older and all older adults:

- All children from 6 months through 18 years of age.
- Anyone 50 years of age or older.

Anyone who is **at risk of complications from influenza, or more likely to require medical care:**

- Women who will be **pregnant** during influenza season.
- Anyone with **long-term health problems** with:
 - heart disease
 - kidney disease
 - liver disease
 - lung disease
 - metabolic disease, such as diabetes
 - asthma
 - anemia, and other blood disorders
- Anyone with a **weakened immune system** due to:
 - HIV/AIDS or other diseases affecting the immune system
 - long-term treatment with drugs such as steroids
 - cancer treatment with x-rays or drugs
- Anyone with certain **muscle or nerve disorders** (such as seizure disorders or cerebral palsy) that can lead to breathing or swallowing problems.
- Anyone 6 months through 18 years of age on **long-term aspirin treatment** (they could develop Reye Syndrome if they got influenza).
- **Residents of nursing homes** and other **chronic-care facilities.**

Anyone who lives with or cares for people at high risk for influenza-related complications:

- **Health care providers.**
- **Household contacts and caregivers of children** from birth up to 5 years of age.
- **Household contacts and caregivers of**
 - people 50 years and older, or
 - anyone with medical conditions that put them at higher risk for severe complications from influenza.

Health care providers may also recommend a yearly influenza vaccination for:

- People who provide **essential community services.**
- People living in **dormitories, correctional facilities,** or under other **crowded conditions,** to prevent outbreaks.
- People at high risk of influenza complications who **travel** to the Southern hemisphere between April and September, or to the tropics or in organized tourist groups at any time.

4 When should I get influenza vaccine?

You can get the vaccine as soon as it is available, usually in the fall, and for as long as illness is occurring in your community. Influenza can occur any time from November through May, but it most often peaks in January or February. Getting vaccinated in December, or even later, will still be beneficial in most years.

Most people need one dose of influenza vaccine each year.

Children younger than 9 years of age getting influenza vaccine for the first time – or who got influenza vaccine for the first time last season but got only one dose – should get 2 doses, at least 4 weeks apart, to be protected.

Influenza vaccine may be given at the same time as other vaccines, including pneumococcal vaccine.

5 Some people should talk with a doctor before getting influenza vaccine

Some people should not get inactivated influenza vaccine or should wait before getting it.

- Tell your doctor if you have any **severe** (life-threatening) allergies. Allergic reactions to influenza vaccine are rare.
 - Influenza vaccine virus is grown in eggs. People with a severe egg allergy should not get the vaccine.
 - A severe allergy to any vaccine component is also a reason to not get the vaccine.
 - If you have had a severe reaction after a previous dose of influenza vaccine, tell your doctor.
- Tell your doctor if you ever had Guillain-Barré Syndrome (a severe paralytic illness, also called GBS). You may be able to get the vaccine, but your doctor should help you make the decision.
- People who are moderately or severely ill should usually wait until they recover before getting flu vaccine. If you are ill, talk to your doctor or nurse about whether to reschedule the vaccination. People with a mild illness can usually get the vaccine.

6 What are the risks from inactivated influenza vaccine?

A vaccine, like any medicine, could possibly cause serious problems, such as severe allergic reactions. The risk of a vaccine causing serious harm, or death, is extremely small.

Serious problems from influenza vaccine are very rare. The viruses in inactivated influenza vaccine have been killed, so you cannot get influenza from the vaccine.

Mild problems:

- soreness, redness, or swelling where the shot was given
- hoarseness; sore, red or itchy eyes; cough
- fever • aches

If these problems occur, they usually begin soon after the shot and last 1-2 days.

Severe problems:

- Life-threatening allergic reactions from vaccines are very rare. If they do occur, it is usually within a few minutes to a few hours after the shot.
- In 1976, a type of influenza (swine flu) vaccine was associated with Guillain-Barré Syndrome (GBS). Since then, flu vaccines have not been clearly linked to GBS. However, if there is a risk of GBS from current flu vaccines, it would be no more than 1 or 2 cases per million people vaccinated. This is much lower than the risk of severe influenza, which can be prevented by vaccination.

7 What if there is a severe reaction?

What should I look for?

Any unusual condition, such as a high fever or behavior changes. Signs of a severe allergic reaction can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat or dizziness.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** the doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your provider to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form. Or you can file this report through the VAERS website at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS does not provide medical advice.

8 The National Vaccine Injury Compensation Program

A federal program exists to help pay for the care of anyone who has a serious reaction to a vaccine.

For more information about the National Vaccine Injury Compensation Program, call **1-800-338-2382**, or visit their website at www.hrsa.gov/vaccinecompensation.

9 How can I learn more?

- Ask your provider. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/flu



DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION



Vaccine Information Statement (Interim)
Inactivated Influenza Vaccine (8/11/09) 42 U.S.C. §300aa-26